# Conceptualizing Psychopathy Using the CAPP Formulation Over Numbers



Stephen D. Hart, PhD



# Clinical Questions

- What is the nature and severity of the patient's psychopathic symptomatology?
- Does the patient currently suffer from psychopathy?

## Clinical Questions

- Was this patient's past violence due to psychopathy?
- Has the patient's violence risk changed due to changes in psychopathic symptomatology?

## Clinical Questions

- What should be the focus of treatment for psychopathy for this patient?
- What type of treatment for psychopathy is most appropriate for this patient?

## Formulation: PCL-R

PCL-R TOTAL SCORE = 32

#### Problems: PCL-R

- Its content is not ideal
  - Some important features of psychopathy are not captured well or even excluded
  - Some features of psychopathy of questionable importance are included

### Problems: PCL-R


- Its format makes it insensitive to change
  - Focus is on lifetime prevalence
  - Many item definitions are anchored in low-frequency, situation-specific behaviors

### Problems: PCL-R

- Its item ratings are psychometrically sound but clinically irrelevant
  - Rating format confounds nature and severity of symptoms
  - Adding ratings to create scores assumes the items are interchangeable


#### CAPP

- A conceptual model intended to provide a definition of psychopathy that is
  - Explicit
  - Comprehensive
  - Lexical




#### CAPP-IRS

- A rating scale based on the CAPP that permits
  - Ratings of overall symptom severity, plus separate ratings of trait extremity and functional impairment
  - Flexible timeframe for ratings



## 3 Goals of Treatment



### Contact Information

SFU

Department of Psychology Simon Fraser University Burnaby, Canada



Faculty of Psychology University of Bergen Bergen, Norway

hart@sfu.ca

http://www.sfu.ca/psyc/faculty/hart/